



## In situ growth of individual graphene layers

**Kling, Jens; Hansen, Thomas Willum; Wagner, Jakob Birkedal**

*Published in:*  
Proceedings of IAM Nano 2015

*Publication date:*  
2015

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Kling, J., Hansen, T. W., & Wagner, J. B. (2015). In situ growth of individual graphene layers. In *Proceedings of IAM Nano 2015*

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## ***In situ* growth of individual graphene layers**

Jens Kling, Thomas W. Hansen, Jakob B. Wagner

*DTU Cen, Lyngby, Denmark*

Chemical vapor deposition (CVD) is a relatively easy and cheap method for large scale synthesis of graphene. Commonly used catalysts are copper or nickel [1], [2] foils or wafers. However, the understanding and control of such growth processes are still in their infancy.

We present *in situ* transmission electron microscopy (TEM) experiments in a FEI Titan 80-300 Environmental TEM (ETEM) for studying the growth of layered carbon materials on these catalysts. The ETEM allows imaging with controlled gas environments around the sample up to a few mbar. In combination with a MEMS-based heating holder, growth of layered carbon materials is systematically studied at the atomic level using various gas pressures and growth temperature.

The growth of few layer graphene from  $C_2H_2$  on a Ni catalyst is shown in Fig. 1. NiO particles in the size range up to a few hundred nm are reduced in the microscope under  $H_2$  at 500-600°C in order to form a catalytically active Ni surface. Introducing  $C_2H_2$  at about 650°C leads to growth of layered carbon. By following the appearance of carbon layers, the growth rate dependence on various parameters can be determined directly from the ETEM observations.

These results should help to better understand the growth mechanism and help to control and optimize the process.

### **References:**

- [1] X. Li, W. Cai, J. An, S. Kim, J. Nah, D. Yang, R. Piner, A. Velamakanni, I. Jung, E. Tutuc, S. K. Banerjee, L. Colombo, and R. S. Ruoff, *Science* **324**, 1312 (2009).
- [2] X. Li, W. Cai, L. Colombo, and R. S. Ruoff, *Nano Lett.* **9**, 4268 (2009).

**Images:**





Fig. 1: *Image series with 0.61s between the images. The in-plane growth of the carbon layers is easily observed (see arrows).*